lan Bookman, MD Gastroenterology

Lecturer, University of Toronto

You have been scheduled for a COLONOSCOPY on	
CHECK-IN T	IME: PROCEDURE TIME:
S	atient Registration, Ground Floor, Tranquility Garden Entrance St. Joseph's Health Centre, 30 The Queensway For questions or directions, visit www.stjoe.on.ca or call 416-530-6156.
CANC	consible adult to escort you home, OR ELSE YOUR PROCEDURE WILL BE ELLED. You will NOT be allowed to drive or work for the rest of that day. The Health Card
Cancelation F	ees: \$250 if less than 48 hours notice \$25 if less than 2 weeks notice
Preparation In 1. Medic	nstructions: ations: Continue all medications, including the day of the test, except for:
	Coumadin – hold for 3 days Aspirin – hold for 7 days Plavix – hold for 7 days - you MUST confirm with your GP or cardiologist if it is OK for you to stop these medications
d.	Insulin – if short-acting, hold morning dose – if 30/70 or long-acting, take half morning dose
e. f	Glyburide – hold morning dose Iron Supplements – hold for 7 days

2. **Purchase:** 1 Box of PICO-SALAX (contains 2 packets) and 1 Bottle of Mag Citrate from *any* pharmacy. You do NOT need a prescription for them.

One Day Before The Colonoscopy:

- 3. You are ONLY allowed to eat or drink the following clear fluids: coffee and tea (NO milk or cream), clear soup (no noodles or solid contents), clear juices (apple, white grape, white cranberry), Gatorade, ginger ale, jello (except red). NO dairy products. Ignore the diet directions that come with the Pico-Salax box. Please Do Not Eat Any Solid Food.
- 4. Drink 3 litres of clear fluids throughout the day. The more you drink, the better it works.
- 5. **10AM** Mix the **1st Packet of Pico-Salax** with 1 glass of cold water and drink.
- 6. 4PM Mix the 2nd Packet of Pico-Salax with 1 glass of cold water and drink.
- 7. **8PM** Drink the **Bottle of Mag Citrate.**
- 8. Have nothing to eat or drink after midnight, except for your medications in the morning with a few sips of water.